



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

base of the petals towards their extremity. Indeed, nearest to their base, within the first third of their length, there is a large irregular spot of an intense yellow; about the middle of their length there follows a narrower cross band of red color, vermilion towards the base, intensely pink towards the outside, not reaching the margin of the petals, sometimes dissolved into several separate spots; lastly, beyond the middle of the length of the petals there are three to eight smaller roundish spots of a paler violet pink color. The flowers of *Veronica chamædrys* prove that also gay blue colors are perceived and selected by *Ascia*.—HERMANN MULLER in *Nature*.

CAREX SULLIVANTII.—Mr. E. C. Howe, of Yonkers, West Chester Co., N. Y., writes that he has collected during the present season several specimens of the above *Carex*, and would like to exchange them for some Western Carices, such as *C. Shortiana*, *C. Meadii*, *C. Bebbii*, *C. crus-corvi*, *C. conjuncta*, *C. muricata*, *C. cephaloidea*, or *C. Fraseriana*.

CROSS FERTILIZATION OF BAPTISIA TINCTORIA.—Prof. W. W. Bailey writes in reference to *B. tinctoria* that it is cross-fertilized by humble bees. Their weight on the keel causes a quick and decided lateral deflection of the wings, exposing the andrœcium. A careful study of this mechanism would be very interesting.

DOWNINGIA PULCHELLA.—In a field east of San Jose I saw last June at least five acres completely carpeted with *Downingia pulchella*. The nearly level ground had been sown with wheat which the April flood "drowned out" in the lowest places. In September the same ground will be covered with cocklebur.

The rare *Mentzelia Lindleyi* is abundant near Alum Rock, seven miles east of San Jose.—V. RATTAN, *San Francisco, Cal.*

NEW LOCALITY FOR SULLIVANTIA OHIONIS.—Happening to spend a day in the eastern part of Cass county, Indiana, I found on the limestone bluffs overhanging Pipe Creek, just before its junction with the Wabash River, *Sullivantia Ohionis* in abundance. The general conformation of the country and the relative situation of *Sullivantia*, are almost exact counterparts of the station in Jefferson Co., Indiana, with the single exception that the bluffs are not nearly so high. The exposure and character of the soil seem to be identical.—M. S. COULTER, *Logansport, Ind.*

SCIENCE—A Weekly Record of Scientific Progress. Illustrated. We have received the initial number of the above journal, which claims to "occupy a field in periodical literature hitherto unoccupied," "and the only first class weekly Journal in the United States devoted to science, recognized by scientists as their medium of communication." Furthermore, all desiring to keep "au courant", or rather